



520 Rec'd PCT/PTO 12 OCT 1999

Attorney Docket No. FCCC-96-11

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Tsichlis, et. al.

Serial No.: 09/202,549

Group Art Unit No.: Not Yet Assigned

Filed: December 16, 1998

Examiner: Not Yet Assigned

For: Nucleic Acid Molecule For Enhancing Gene Expression

Assistant Commissioner of Patents
Washington, D.C. 20231

**RESPONSE TO NOTIFICATION OF
MISSING REQUIREMENTS UNDER 35 U.S.C. § 371**

Sir:

In response to the "Notification of Missing Requirements Under 35 U.S.C. § 371" dated August 12, 1999, enclosed herewith for filing is the Combined Declaration and Power of Attorney, executed by the inventors.

This response, with the concurrently filed Petition for One Month Extension of Time, is believed to be timely on October 12, 1999.

The fee to cover the surcharge (\$65.00) and one-month extension of time (\$55.00) is enclosed herewith.


10/21/1999 PVOLPE 00000113 09202549

01 FC:215

55.00 OP

Date: October 12, 1999


Respectfully submitted,


William J. McNichol, Jr.
Registration No. 31,179
Attorney for Applicants

REED SMITH SHAW & McCLAY
2500 One Liberty Place
1650 Market Street

Repln. Ref: 10/21/1999 PVOLPE 00000113 09202549
DAH:180586
FC: 704

Att: William J. McNichol, Jr., Esq.
(215 241-7950)



2500 One Liberty Place
1650 Market Street
Philadelphia, Pennsylvania 19103-7301
Phone: 215-851-8100
Fax: 215-851-1420
October 12, 1999

RE	Serial No.:	09/202,549
	Applicant:	Tsichlis et. al.
	Title:	Nucleic Acid Molecule For Enhancing Gene Expression
	Filing Date:	December 16, 1998
	Docket No.:	FCFC-96-11

1. *Chlorophyll a* and *Chlorophyll b* were determined by the method of Lichtenthaler and Whistler (1973). The total chlorophyll content was determined by the method of Arar and Cook (1980). The carotenoid content was determined by the method of Lichtenthaler and Whistler (1973). The total carotenoid content was determined by the method of Arar and Cook (1980). The total protein content was determined by the method of Lowry et al. (1951). The total lipid content was determined by the method of Bligh and Dyer (1959). The total carbohydrate content was determined by the method of Dubois and Gilles (1950). The total nucleic acid content was determined by the method of Burton (1956). The total ash content was determined by the method of AOAC (1990). The total moisture content was determined by the method of AOAC (1990). The total dry matter content was determined by the method of AOAC (1990). The total organic acid content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenol content was determined by the method of AOAC (1990). The total terpenoid content was determined by the method of AOAC (1990). The total steroid content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990). The total alkaloid content was determined by the method of AOAC (1990). The total saponin content was determined by the method of AOAC (1990). The total tannin content was determined by the method of AOAC (1990). The total flavonoid content was determined by the method of AOAC (1990). The total phenol content was determined by the method of AOAC (1990). The total terpenoid content was determined by the method of AOAC (1990). The total steroid content was determined by the method of AOAC (1990). The total glycoside content was determined by the method of AOAC (1990).